



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\therefore d(a/x) = 2k(n+1)d\delta.$$

$$\therefore dp = 2k(n+1)ag\delta d\delta, p = agk(n+1)\delta^2.$$

Let  $l$  = height of a homogenous atmosphere of density  $\rho$  exerting a pressure  $p_0$ .

$$\therefore p_0 = g\rho l.$$

$$\therefore \frac{p}{p_0} = (n+1) \frac{a}{l} \cdot \frac{k\delta^2}{\rho}, \text{ when } p = p_0, \delta = \rho.$$

$$\therefore 1 = (n+1) \frac{a}{l} \cdot k\rho, \text{ or } n = \frac{l/a}{k\rho} - 1.$$

$$a = 6366738 \text{ meters, } k\rho = .000147192.$$

$$l = .76 \text{ (as many times as mercury is as heavy as air)} = 7993.15 \text{ meters.}$$

$$\therefore l/a = .00125545.$$

$$\therefore n = \frac{.00125545}{.000147192} - 1 = 7.5, \text{ a value of } n \text{ too large.}$$

---

## DEPARTMENTS.

---

### SOLUTIONS OF PROBLEMS.

---

#### ARITHMETIC.

---

146. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

If the driving-wheels of Locomotive No. 200 on the Pennsylvania Railroad,  $m=7$  feet in diameter, turn  $n=20$  times in  $p=3$  seconds, and lose  $r=12\%$  of their forward motion by slipping on the smooth steel rails, at what rate per hour is the locomotive moving over the rails?

Solution by W. P. WEBBER, Mississippi Normal College, Houston, Miss.

$m\pi$  = circumference of wheel,  $mn\pi$  = number of times the circumference is applied to the rail, and is therefore the distance the engine travels without slipping in  $p$  seconds.

Hence,  $\frac{mn\pi}{p}$  = distance engine travels in 1 second without slipping, and  $\frac{mn(100-r)\pi}{100p}$  = actual distance engine travels in 1 second, since it slips back  $r\%$  of the distance it travels.

Substituting numbers for the letters, we have for the distance the engine travels in 1 second,  $\frac{7 \times 20(100-12)\pi}{100 \times 3}$  feet.

Hence, the rate of the engine in miles per hour is

$$\frac{7 \times 20 \times 88 \times \pi \times 3600}{100 \times 3 \times 5280} = 28\pi \text{ miles.}$$

---

**ALGEBRA.**

---

123. Proposed by **ELMER SCHUYLER**, B. Sc., Professor of German and Mathematics, Boys' High School, Reading, Pa.

$$\left(\frac{1+x}{1-x}\right)^{\frac{1}{4}} + \sqrt{\frac{1-a}{1+a}} \sqrt[4]{\frac{1-x}{1+x}} = 2 \sqrt[4]{\frac{1-a^2}{(1+a)^2}}, \text{ and } \sqrt{a^2-x^2} + x\sqrt{a^2-1} = a^2\sqrt{1-x^2}.$$

[Haddon.]

Solution by **JOHN A. VAN GROOS**, Fellow of Mathematics, University of Oregon, Eugene; Ore.; **R. L. MOORE**, Student in University of Texas, 2206 San Marcos Street, Austin, Tex.; and **G. B. M. ZERR**, A. M., Ph.D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

$$(1). \text{ Multiply } \left(\frac{1+x}{1-x}\right)^{\frac{1}{4}} + \sqrt{\frac{1-a}{1+a}} \sqrt[4]{\frac{1-x}{1+x}} = 2 \sqrt[4]{\frac{1-a^2}{(1+a)^2}} = 2 \sqrt[4]{\frac{1-a}{1+a}} \text{ through by}$$

$$\left(\frac{1+x}{1-x}\right)^{\frac{1}{4}} \cdot \left(\frac{1+x}{1-x}\right)^{\frac{1}{4}} - 2 \left(\frac{1-a}{1+a}\right)^{\frac{1}{4}} \left(\frac{1+x}{1-x}\right)^{\frac{1}{4}} + \left(\frac{1-a}{1+a}\right)^{\frac{1}{4}} = 0.$$

$$\therefore \left[ \left(\frac{1+x}{1-x}\right)^{\frac{1}{2}} - \left(\frac{1-a}{1+a}\right) \right]^2 = 0. \quad \therefore \frac{1+x}{1-x} = \frac{1-a}{1+a}.$$

$$\therefore x = -a.$$

$$(2). \sqrt{a^2-x^2} = a^2\sqrt{1-x^2} - x\sqrt{a^2-1}.$$

$$a^2-x^2 = a^4 - a^4x^2 - 2a^2x\sqrt{(1-x^2)(a^2-1)} + a^2x^2 - x^2.$$

$$\therefore 2a^2x\sqrt{(1-x^2)(a^2-1)} = a^2(1-x^2)(a^2-1).$$

$$\therefore \{2x - \sqrt{[(1-x^2)(a^2-1)]}\} \sqrt{[(1-x^2)(a^2-1)]} = 0.$$

$$\therefore 4x^2 = (1-x^2)(a^2-1) \text{ or } x^2 = 1.$$

$$\therefore x = -1, +1, \text{ or } \pm \sqrt{\frac{a^2-1}{3+a^2}}.$$

The values  $x = -1$  and  $\pm \sqrt{\frac{a^2-1}{3+a^2}}$  are the values satisfying the equation as given.

124. Proposed by **J. SCHEFFER**, A. M., Hagerstown, Md.

A certain quantity of alcohol diluted with water so that in one liter there are  $c$  liters of pure alcohol, is mixed  $n$  times successively with  $p$  times the quantity of alcohol diluted so that 1 liter contains a liter of pure alcohol. How much pure alcohol does one liter of the  $n$ th mixture contain?